

Numerical and Theoretical Investigations of North Pacific Subtropical Mode Water with Implications to Pacific Climate Variability

Xujing Jia Davis

Graduate School of Oceanography, University of Rhode Island

Lewis M. Rothstein

Graduate School of Oceanography, University of Rhode Island

William K. Dewar

Department of Oceanography, Florida State University,


Dimitris Menemenlis

Jet Propulsion Laboratory, California Institute of Technology



North Pacific Subtropical Mode Water (STMW)

Schematic current patterns in western North Pacific



Location: forms and resides south of Kuroshio Extension (KE)

Features:

- weakly stratified, low PV
- upper 500 m of the ocean water column
- inhabits thermostads between 16 and 19°C
- salinity range of 34.65-34.8psu
- potential density values of 24.8-25.7 kg/m³

(Masuzawa, 1969; Suga et al., 1990; Eitarou et al., 2004)

Questions

- STMW has known seasonal variability, but what is the variability of STMW on longer time scales?
- What is the relationship (if any) between low frequency STMW and established climate patterns in the Pacific?
- Dynamics behind it?

Model and Simulation Descriptions

MITgcm : 3D, z level, primitive equation OGCM (Marshall, 1997)

➤ ECCO2 (Cube 37) global-ocean and sea-ice simulation :

-28-year spin-up prior to its initial January 1992 conditions, carried out by cycling through the 1992-2000 NCEP forcing converted to fluxes using model SST and the Large and Pond bulk formulae (*Large et al, 1995, Menemenlis, 2005*)

➤ θ , S, u, v

- Resolution: horizontal resolution: 1/6 lat x 1/6 lon; vertically, from surface to ~6km, 10 m resolution above 100 m and stretched to 95 m around 1000 m

- temporal coverage: 1992, Jan – 2006 Mar (171 months)

-Output hasn't been constrained by oceanic and seaice data yet


STMW definition

Definition in Cube37 simulation

- PV is less or equal to $2 \times 10^{-10} m^{-1} s^{-1}$

$$PV = -\frac{f}{\rho} \frac{\partial \sigma_{\theta}}{\partial z}$$


- potential density between 24.5~25.3 kg/m³
- region of 130E~ 200E, 20N~ 40N and east of islands of Japan




Defined STMW region

Compare with Observation

MITgcm, May 2004




KESS, late May 2004




After Rainville, et al., 2007

Temporal variability: 3-D averaged features of STMW




During 1999/2000, cooler, fresher, lower PV, lighter, thinner, shallower STMW

Temporal variability: STMW volume



Dominant signal for the STMW variability



Annual cycle is the most significant

Seasonality of STMW in climatological fields in MITgcm


Take θ as an example:

$$\theta_{month}(lon, lat) = \frac{1}{14} \sum_{year=1992}^{2005} \theta_{year, month}(lon, lat) \quad (month=1, 2, \dots, 12)$$

Where $\theta_{month}(lon, lat)$ is the calculated climatological field


Seasonality in climatological fields

-meridional cross section




Seasonality in climatological fields

-meridional cross section




Seasonality in climatological fields

-meridional cross section




Seasonality in climatological fields

-meridional cross section




Results: Seasonality in climatological fields

-meridional cross section




Seasonality in climatological fields

-meridional cross section



Results: Seasonality in climatological fields

-three time periods





Period I: STMW formation (Nov~Mar)

Period II: STMW isolation (Mar~Jun)

Period III: STMW dissipation (Jun~Nov)



Interannual variability of STMW

STMW interannual variability




STMW variability and its relation to Pacific climate variation

The Pacific Decadal Oscillation (PDO)



www.jisao.washington.edu/pdo/

STMW Variability & PDO




STMW variability is highly correlated with PDO index
Co=0.69, significant value=0.1 with 95% level of confidence 22


Connection between STMW & PDO:
large scale atmospheric variations

PDO Index

The 1st EOF time coefficient of the SST north of 20 N in Pacific



Connection between STMW & PDO: large scale atmospheric variations



Summary (MITgcm)

- The dominant temporal pattern of STMW is seasonality, the annual cycle can be divided into formation, isolation and dissipation periods that correspond to distinct stages of STMW evolution
- An interannual signal is clearly seen in STMW variability as well, this lower frequency signal shows significant correlation with PDO index
- This likely results from the variations in the large scale atmospheric forcing: wind stress and air-sea heat flux

Theoretical frameworks and possible mechanisms of STMW variability

- Following Dewar et al 2005, a modified LPS framework may be established to describe STMW and its connections to large scale ocean-atmospheric circulation
- A PGOM (Samelson & Vallis, 1997) numerically approximates the solutions to this framework and describes STMW characteristics.
- PGOM experiments demonstrate that the interannual variability observed in the Cube 37 simulations can be driven by variations in the large scale air-sea heat flux and wind stress patterns seen in the NCEP reanalysis.

Thank you!


Acknowledgements

- *Lewis Rothstein (URI)*
- *William Dewar (FSU)*
- *Dimitris Menemenlis (JPL)*

- *Roger Samelson (OSU)*
- *Geoffrey Vallis (GFDL)*
- *Young-Oh Kwon (WHOI)*

Seasonality in climatological fields

-meridional cross section



Summary (PGOM)

- PGOM representation of the modified LPS framework produces a distinct analog to STMW
- Within this model/framework, Ekman pumping is necessary for the existence and maintenance of STMW
- Quasi-realistic time varying atmospheric forcing experiments show variable large scale wind stress (Ekman pumping) and air sea heat fluxes can separately generate seasonal and interannual variability in STMW